

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 898 900 A2

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
03.03.1999 Bulletin 1999/09

(51) Int. Cl.⁵: A23L 1/0528, A23L 1/308

(21) Application number: 98201018.3

(22) Date of filing: 31.03.1998

(84) Designated Contracting States:
AT BE CH DE DK ES FI FR GB GR IE IT LI LU MC
NL PT SE
Designated Extension States:
AL LT LV MK RO SI

(72) Inventors:
• Jaussan, Véronique
1110 Morges (CH)
• Roessle, Claudia
1110 Morges (CH)
• Schwelzer, Thomas
1052 Le Mont-Sur-Lausanne (CH)
• Bourguignon, Michel
14740 Lasson (FR)

(30) Priority: 23.06.1997 EP 97201915

(71) Applicant:
SOCIÉTÉ DES PRODUITS NESTLÉ S.A.
1800 Vevey (CH)

(54) Composition and method for providing nutrition to diabetics

(57) A nutritional composition for diabetics. The composition contains a protein source, a lipid source, and a carbohydrate source. The composition includes a fibre mixture containing a viscous soluble fibre and inulin, a hydrolysate of inulin, or both. The fibre mixture may also contain insoluble fibre. When in liquid form, the nutritional composition has a low viscosity and excellent tube flow rates.

EP 0 898 900 A2

Description

Field of the Invention

- 5 [0001] This invention relates to a nutritional composition suitable for providing nutrition to diabetic patients. The invention also relates to a method for providing nutrition to diabetic patients and to a method for the nutritional management of diabetic symptoms in patients.

Background of the Invention

- 10 [0002] Diabetes is a general term for a group of metabolic disorders which are characterised by the inability to properly metabolise glucose. This inability is due either to a deficiency of the hormone insulin or to a resistance to the action of insulin. In either case, if untreated, the inability leads to hyperglycaemia and its complications of morbidity and mortality.
- 15 [0003] Diabetes is usually classified into three clinical classes; diabetes mellitus, gestational diabetes, and impaired glucose tolerance or glucose intolerance. Diabetes mellitus is generally broken down into two types; type I diabetes mellitus and type II diabetes mellitus. Type I diabetes mellitus occurs in individuals who produce little or no insulin and constitutes less than about 10% of the diabetic population. The onset of type I diabetes mellitus usually manifests itself during youth. Type II diabetes mellitus, or non-insulin dependent diabetes mellitus, usually develops after the age of
- 20 about 30 years and constitutes more than about 90% of the diabetic population. Impaired glucose tolerance, in the clinical setting, often occurs as stress-induced hyperglycaemia.
- [0004] Often, especially in severe cases, diabetes is treated by the administration of exogenous insulin or antihyperglycaemic agents. However, nutritional management of diabetes is possible and the American Diabetes Association has published guidelines for the nutritional therapy of diabetes. These guidelines suggest that diabetic patients should
- 25 consume about 20 to 25 g of dietary fibre daily.
- [0005] Dietary fibres may be classified as soluble and insoluble. Insoluble fibres appear to have little influence on glycaemic levels. However, the incorporation of soluble fibres into foods is known to reduce postprandial glycaemia in diabetics (Anderson, J.W. and Akanji, A.O.; 1993; "Treatment of Diabetes with High Fiber Diets", CRC Handbook of Dietary Fiber in Human Nutrition, CRC Press Inc, 2nd Edition, pages 443 to 470). Examples of soluble fibres which are believed
- 30 to have this property are guar gum, pectin, xanthan gum, and β -glucan. This property makes soluble fibres ideal candidates for incorporation into foods for diabetics.
- [0006] However, to date, soluble fibres have not been widely used in food for diabetics. A common problem is that many soluble fibres are not very palatable. Also, in the field of clinical nutrition, the incorporation of soluble fibres into enterally administered compositions is extremely difficult since soluble fibres are thickening agents and greatly increase viscosity. Consequently compositions which contain sufficient soluble fibre to significantly reduce postprandial glycaemia are usually too thick and viscous for enteral feeding; especially for patients requiring tube-feeding.
- 35 [0007] One nutritional composition which offers a solution to the problem is described in US patent 5,292,723. The nutritional composition described in this patent has a carbohydrate component made up of glucose polymers, modified starch, and soluble fibre. Pectin is suggested as a suitable soluble fibre. The nutritional composition has a viscosity of less than 0.05 kg/ms when measured at about 20°C. Further the nutrition composition, when consumed, results in lower glycaemic response in patients as compared to glucose of the same energy content.
- [0008] However there is still a need for a nutritional composition which is suitable for diabetic patients and which has good flow characteristics.

Summary of the invention

- 45 [0009] Accordingly, in one aspect, this invention provides a nutritional composition for diabetic patients, the composition containing a protein source, a lipid source, and a carbohydrate source, the carbohydrate source including a fibre mixture comprising a viscous soluble fibre and inulin or a hydrolysate of inulin, or both.
- 50 [0010] It is surprisingly found that the use of a fibre mixture of a viscous soluble fibre and inulin, a hydrolysate of inulin, or both, results in an adequately reduced glycaemic response while retaining reasonably low viscosity. Therefore, due to the excellent flow properties, the nutritional composition is ideally suited to be tube-fed to patients. Further, the use of inulin or its hydrolysates provides a substrate for lactic acid bacteria in the gastro-intestinal tract; leading to beneficial effects in the general health of the patient.
- 55 [0011] The nutritional composition may be in liquid form or in the form of a soluble powder which is reconstitutable in an aqueous liquid to provide a liquid nutritional composition.
- [0012] Preferably, the protein source provides about 10% to about 20% of energy, the lipid source provides about 30% to about 50% of energy, and the carbohydrate source provides about 35% to about 55% of energy.

[0013] In another aspect, this invention provides a method for the nutritional management of the symptom of diabetes, the method comprising orally administering to a patient an effective amount of a liquid nutritional composition for diabetic patients, the composition comprising a protein source, a lipid source, a carbohydrate source, and a fibre mixture including a viscous soluble fibre and inulin, a hydrolysate of inulin, or both.

Detailed Description of the Preferred Embodiments

[0014] Embodiments of the invention are now described by way of example only. This invention provides a nutritional composition for diabetic patients. The composition contains a protein source, a lipid source, and a carbohydrate source.

The carbohydrate source includes a fibre mixture comprising a viscous soluble fibre and inulin. In this specification, the term "soluble fibre" means those dietary fibres which are characterised as soluble using the method of Prosky et al; 1988; *J. Assoc. Off. Anal. Chem.* 70, 5, 1017. This is the official method of the Association of Official Analytical Chemists. The term "viscous soluble fibre" means a soluble fibre which is able to increase the viscosity of the contents of the stomach and the small intestine and slow gastric emptying rates.

[0015] The viscous soluble fibre may be any suitable soluble fibre which is able to increase the viscosity of the contents of the stomach and the small intestine. Examples of suitable soluble fibres are gums such as guar gum, xanthan gum, and gum arabic, pectin, and β -glucan, or mixtures of these. Pectin and gum arabic are particularly preferred.

[0016] The inulin may be provided in the form of a natural extract which is suitable for human consumption. Extracts from chicory are particularly suitable. The extract preferably contains at least 80% by weight of inulin or a hydrolysate of inulin; more preferably at least 90% by weight of inulin or a hydrolysate of inulin. The inulin preferably has a degree of polymerisation of at least about 8; for example about 10 to about 25. Suitable inulin extracts may be obtained from Orafit SA of Trelmont 3300, Belgium under the trade mark "Rafiline". For example, the inulin may be provided in the form of Rafiline®ST which is a fine white powder which contains about 90 to about 94% by weight of inulin, up to about 4% by weight of glucose and fructose, and about 4 to 9% by weight of sucrose. The average degree of polymerisation of the inulin is about 10 to about 12. The inulin may also be in the form of inulin hydrolysates or mixtures of inulin and inulin hydrolysates. Inulin hydrolysates are commonly known as fructooligosaccharides or FOS.

[0017] Inulin and its hydrolysates are reported to promote the growth of bifidobacteria in the gastro-intestinal tract and, in certain circumstances prevent or decrease the growth of pathogens such as *Clostridia*. Further, promoting the growth of bifidobacteria is reported to have various other beneficial effects. Moreover, inulin and its hydrolysates may reduce blood glucose levels.

[0018] The fibre mixture may also contain a source of insoluble dietary fibre. Suitable sources of insoluble dietary fibres are hull fibres from legumes and grains; for example pea hull fibre, oat hull fibre, barley hull fibre, and soy hull fibre. Pea hull fibre is especially preferred. However any suitable source of insoluble dietary fibre may be used. These sources may also contain some soluble fibre.

[0019] The ratio of soluble fibre, including inulin, to insoluble fibre is preferably about 1:3 to about 3:1; more preferably about 1:1 to about 2:1. For example, ratio of soluble fibre, including inulin, to insoluble fibre is preferably about 2:1. Further, the fibre mixture may be present in an amount of about 1.0 g/100ml to about 2.0 g/100ml; for example about 1.3 g/100ml to about 1.7 g/100ml; for example about 1.5 g/100 ml. In a preferred example, the fibre mixture comprises about 0.5 g/100 ml inulin, about 0.5 g/100 ml pectin or gum arabic, and about 0.5 g/100 ml outer pea fibre.

[0020] The protein source is preferably a high quality protein source; for example milk protein, whey protein, casein protein, or soy protein, or mixtures of these proteins. The protein source may be in the form of intact protein or may be hydrolyzed. Other protein sources such as rice, pea and oat protein, or mixtures thereof, may also be used. Further, if desired, the protein source may include free amino acids.

[0021] The protein source preferably provides about 10% to about 20% of the energy of the composition. For example, the protein source may provide about 12% to about 18% of the energy of the composition; preferably about 15% of the energy of the composition.

[0022] The carbohydrate source may be any suitable carbohydrate or carbohydrate mixtures. For example, the carbohydrate source may be maltodextrin, modified starch, amylose starch, topioca starch, corn starch, or fructose, or mixtures thereof. Modified starch is preferred; especially modified topioca and corn starches. If the carbohydrate source includes maltodextrin, maltodextrin with a low DE is preferred; for example maltodextrin with a DE of 3 or less. Preferably the composition is low in or free from mono- and di-saccharides such as fructose and other exchange sugars, and lactose. For example the composition contains less than about 3 g/l of lactose.

[0023] The carbohydrate source, including the viscous soluble fibre and inulin, provides about 35% to about 55% of the energy of the composition; preferably about 40% to about 50% of the energy. For example, the carbohydrate source may provide about 45% of the energy of the composition. The amount of energy provided to the patient by the viscous soluble fibre and inulin is very small.

[0024] The lipid source is preferably rich in monounsaturated fatty acids; for example monounsaturated fatty acids may provide at least 50% of energy of the lipid source. Preferably, monounsaturated fat acids provide about 25% to

about 35% of the energy of the composition; for example about 29 to 30%. The lipid source may contain polyunsaturated fatty acids (omega-3 and omega-6 fatty acids); preferably these polyunsaturated fatty acids provide up to about 10% of the energy of the composition. For example these polyunsaturated fatty acids may provide about 3% to about 10% of the energy of the composition. The lipid profile of the enteral composition is preferably designed to have a polyunsaturated fatty acid omega-6 (n-6) to omega-3 (n-3) ratio of about 4:1 to about 10:1. Saturated fatty acids preferably provide less than 10% of the energy of the composition; for example less than about 7%.

[0025] The lipid source may provide about 30% to about 50% of the energy of the composition; preferably about 35% to about 45%. For example, the lipid source may provide about 40% of the energy of the composition.

[0026] If desired, the lipid source may include medium chain triglycerides. For example, medium chain triglycerides may make up about 10% to about 50% by weight of the lipid source.

[0027] Suitable sources of monounsaturated fatty acids are olive oil, sunflower oil rich in oleic acid, rapeseed oil rich in oleic acid, hazelnut oil, safflower oil, and the like. If medium chain triglycerides are included in the lipid source, fractionated coconut oils are a suitable source of medium chain triglycerides. A mixture of sunflower oil, rapeseed oil and olive oil is preferred.

[0028] The enteral composition preferably includes a complete vitamin and mineral profile. For example, sufficient vitamins and minerals may be provided to supply about 75% to about 250% of the recommended daily allowance of the vitamins and minerals per 1000 calories of the nutritional composition.

[0029] The nutritional composition conveniently has an osmolality of about 180 mOsm/l to about 300 mOsm/l; for example about 190 mOsm/l to about 210 mOsm/l.

[0030] The viscosity of the nutritional composition, when measured at room temperature, is preferably less than about 0.04 kg/ms; especially less than about 0.035 kg/ms. For example, the viscosity of the nutritional composition, when measured at room temperature, may be about 0.015 to about 0.03 kg/ms. Further, the flow rate of the nutritional composition through a standard feeding tube is preferably less than about 150 minutes/l, for example less than about 100 minutes/l.

[0031] The energy density of the nutritional composition is preferably about 700 kcal/l to about 1500 kcal/l; for example about 1000 kcal/l.

[0032] The nutritional composition is preferably in the form of a ready-to-use formulation. In this form, the composition may be fed to a patient via a nasogastric tube, jejunum tube or by having the patient drink it. As such, the nutritional composition may be in a variety of forms; for example as a fruit juice-type beverage, a milk shake-type beverage and the like. However, the nutritional composition may be in soluble powder form to be reconstituted prior to use.

[0033] Various flavours, sweeteners and other additives may be present. Artificial sweeteners such as acetosulfame and L-aspartyl based sweeteners may be used; for example aspartame.

[0034] The nutritional composition may be produced as is conventional; for example, by blending together the protein source, the carbohydrate source, and the lipid source. If used, the emulsifiers may be included in the blend. The vitamins and minerals may be added at this point but are usually added later to avoid thermal degradation. Any lipophilic vitamins, emulsifiers and the like may be dissolved into the lipid source prior to blending. Water, preferably water which has been subjected to reverse osmosis, may then be mixed in to form a liquid mixture. The temperature of the water is conveniently about 50°C to about 80°C to aid dispersal of the ingredients. Commercially available liquefiers may be used to form the liquid mixture.

[0035] The liquid mixture may then be thermally treated to reduce bacterial loads. For example, the liquid mixture may be rapidly heated to a temperature in the range of about 80°C to about 110°C for about 5 seconds to about 5 minutes. This may be carried out by steam injection or by heat exchanger; for example a plate heat exchanger.

[0036] The liquid mixture may then be cooled to about 60°C to about 85°C; for example by flash cooling. The liquid mixture is then homogenised; for example in two stages at about 7 MPa to about 40 MPa in the first stage and about 2 MPa to about 14 MPa in the second stage. The homogenised mixture may then be further cooled to add any heat sensitive components; such as vitamins and minerals. The pH and solids content of the homogenised mixture is conveniently standardised at this point.

[0037] For a product in liquid form, the homogenised mixture is preferably aseptically filled into suitable containers. Aseptic filling of the containers may be carried out by pre-heating the homogenised mixture (for example to about 75 to 85°C) and then injecting steam into the homogenised mixture to raise the temperature to about 140 to 160°C; for example at about 150°C. The homogenised mixture may then be cooled, for example by flash cooling, to a temperature of about 75 to 85°C. The homogenised mixture may then be further homogenised, cooled to about room temperature and filled into containers. Suitable apparatus for carrying out aseptic filling of this nature is commercially available.

[0038] For a product in powder form, the homogenised mixture is dried to powder; for example by spray drying. Conventional procedures may be used.

[0039] The nutritional composition may be used as a nutritional support for patients suffering from metabolic anomalies which make them susceptible to hypo- or hyperglycaemia. For example, the nutritional composition may be used as a nutritional support for patients suffering from type I diabetes mellitus, type II diabetes mellitus, or intolerance to glu-

cose. Further, the nutritional composition may be used as a nutritional support for patients who are at risk of a recurrence of hypo- or hyperglycaemia. The nutritional composition may also be used as nutritional support for post operative patients. The nutritional composition is particularly useful for the nutritional management of diabetic symptoms of diabetic patients.

- 5 [0040] The amount of the nutritional composition required to be fed to a patient will vary depending upon factors such as the patient's condition, the patient's body weight, the age of the patient, and whether the nutritional composition is the sole source of nutrition. However the required amount may be readily set by a medical practitioner. In general, sufficient of the nutritional composition is administered to provide the patient with about 1 g protein to about 4.0 g protein per kg of body weight per day. For example, an adult patient may be administered about 1.5 g protein to about 2.0 g protein per kg of body weight per day. Further, sufficient of the nutritional composition is administered to provide the patient with up to about 40 g of dietary fibre (insoluble and soluble) per day; for example about 25 g to about 35 g of dietary fibre per day. If the nutritional formula is used as a supplement to other foods, the amount of the nutritional composition that is administered daily may be decreased accordingly.
- 10 [0041] The nutritional composition may be taken in multiple doses, for example 2 to 5 times, to make up the required daily amount or may taken in a single dose. The nutritional composition may also be fed continuously over a desired period.

Example 1

- 20 [0042] A ready-to-use nutritional composition is prepared. The nutritional composition includes the following components:

25

30

35

40

45

50

55

	Component	Conc. (/100ml)	Energy (%)
5	Protein	3.8 g	15
	Casein/Soy Protein (1:1)		
	Carbohydrate	11.2 g	45
	Maltodextrin (low DE)	1.0 g	
10	Modified Starch	10.2 g	
	Soluble fibre including Inulin	1.0 g	
	Insoluble fibre	0.5 g	
15	Lipids	4.4 g	40
	Rapeseed oil, Sunflower oil, Olive oil, Glyceryl stearate, Soya lecithin		
	Vitamins		
20	Vitamin A	150 IU	
	Vitamin C	10 mg	
	Vitamin D	10 IU	
	Vitamin E	1.0 mg	
25	Vitamin K	3.0 µg	
	Thiamin	0.1 mg	
	Riboflavine	0.12 mg	
	Pantothenic acid	0.50 mg	
30	Vitamin B6	0.14 mg	
	Vitamin B12	0.30 µg	
	Niacin	1.2 mg	
35	Folic acid	18 µg	
	Biotin	10 µg	
	Minerals		
40	Zinc, Iron, Copper, Magnesium, Manganese, Selenium, Iodine, Potassium, Calcium, Phosphorous, Chloride		

[0043] The composition has an osmolality of 210 mOsm/l and an osmolality of 240 mOsm/kg. The viscosity is 0.023 kg/ms and the free flow rate is less than 70 minutes per 500 ml through standard enteral tubing.

Example 2

[0044] Eight healthy volunteers, of both sexes, are used in the study. The volunteers are between 20 and 45 years of age and have fasting blood glucose levels of about 70 to about 110 mg/dl. Any volunteer exhibiting the symptoms of diabetes mellitus type I or II or fructose intolerance are excluded.

[0045] The study is carried out in two stages, each stage comprising two study days separated by a wash out period of at least 7 days. The stages are also separated by a wash out period of at least 7 days. Prior to each study day, each volunteer consumes an evening meal of pizza, salad and an apple. No alcohol is taken. Then, from 10 pm, each volunteer undergoes an overnight fast. In the morning of the study day, an indwelling catheter is placed on each volunteer. A blood sample is then taken. A meal is then consumed within 10 minutes of the taking of the sample. Further blood samples are taken at 15, 30, 45, 60, 90, 120 and 180 minutes after meal intake. The blood glucose for each sample is analysed by the glucose oxidase method using a COBAS analyser (Hoffmann-La Roche). The insulin level for each sample is measured by radioimmuno assay (Pharmacia).

[0046] Blood glucose levels are analysed using a two-way ANOVA. The integrated area under the curve (AUC) is calculated using the method of Wolever *et al*; 1986; *Am. J. Clin. Nutr.*, 43, 167-172. The difference between the AUC of the various curves is evaluated using a Wilcoxon matched pairs test or the Hill Armitage test for cross-over design. The glycemic response is considered as different if the AUC curves are statistically different with $p < 0.05$.

[0047] Three products are used as the meals; the product of example 1, the Sondalis[®] Fiber product of Nestlé Clinical Nutrition, and the Fresubin Diabetes product of Fresenius GmbH. Four hundred 400 ml of the product of example 1, 400 ml of the Sondalis[®] Fiber product, and 444 ml of the Fresubin Diabetes product are consumed in each case to provide a standard energy intake of 400 kcal. All meals contain comparable amounts of minerals and micronutrients.

[0048] On the first study day of stage 1, 4 volunteers are fed the product of example 1 and 4 volunteers are fed the Sondalis[®] Fiber product. The selection of volunteers for any product is random. On the second day of stage 1, the products are reversed. On the first day of stage 2, 4 volunteers are fed the product of example 1 and 4 volunteers are fed the Fresubin Diabetes product. The selection of volunteers for any product is random. On the second day of stage 2, the products are reversed.

[0049] The results are as follows:

15

Time (Minutes)	Plasma Glucose Mean (mmol/l)		
	Product Example 1	Sondalis [®] Fiber Product	Fresubin Diabetes product
-10	4.88	4.85	5.01
15	5.24	5.56	5.36
25	5.21	6.29	6.31
45	4.70	5.28	5.65
60	4.43	4.80	4.60
90	4.35	4.84	4.28
120	4.64	4.22	4.32
180	4.47	4.06	4.47

[0050] The area under the curve results are as follows:

	Product Example 1	Sondalis [®] Fiber Product	Fresubin Diabetes product
Area at 120 minutes (mmol/l)	37	84	49

[0051] The glycemic responses of the product of example 1 and the Fresubin Diabetes product are significantly lower than that of the Sondalis[®] Fiber; about 30 mmol/l and 50 mmol/l respectively after 120 minutes compared to about 80 mmol/l after 120 minutes. Further, for the product of example 1, blood glucose response is much flatter than that of the Sondalis[®] Fiber product. The peak change of the product of example 1 is also less than that of the other products. While the glycemic response of the product of example 1 is lower than that of the Fresubin Diabetes product, the differences are not significant. Therefore the product of example 1 is suitable for use with diabetic patients.

[0052] None of the patients indicated any symptoms of digestive intolerance.

Example 3

[0053] The viscosity and flow rate of the composition of example 1 and the product sold by Fresenius GmbH under the name Fresubin DFN Plus are determined:

[0054] The viscosity of each composition is determined at 25°C using a Contraves Rheomat according to manufacturer's instructions. The composition of example 1 is filled into a sealed, flexible bag and connected to a stand at a standard height. The Fresubin DFN Plus product, which is in a glass container (its original packing) is connected to a

separate stand at the same height. An enteral feeding tube is connected to each container and the time required to deliver 500 ml through the enteral feeding tube is determined. An open reservoir is then connected to each stand and the composition of example 1 poured into one reservoir and the Fresubin DFN Plus product poured into the other. An enteral feeding tube is connected to each reservoir and the time required to deliver 500 ml through the enteral feeding tube is determined.

Property	Composition of example 1	Fresubin DFN Plus product
Viscosity (kg/ms)	0.023	0.035
Free Flow Rate (min/500ml)		
- Original Container	66	158
- Open reservoir	60	129

[0055] The results indicate that the composition of example 1 has a superior flow rate.

Example 4

[0056] A ready-to-use nutritional composition is prepared. The nutritional composition includes the following components:

Component	Conc. (g/100ml)	Energy (%)
Protein	3.8 g	15
Casein/Soy Protein (1:1)		
Carbohydrate	11.2 g	45
Maltodextrin (low DE)	1.0 g	
Modified tapioca and corn starch	10.2 g	
Soluble fibre including inulin	1.0 g	
Insoluble fibre	0.5 g	
Lipids	4.4 g	40
Rapeseed oil, Sunflower oil, Olive oil, Mono- and di-glycerides (E471)		
Vitamins and minerals as in example 1		

[0057] The composition has an osmolality of 190 mOsm/l. The viscosity is about 0.023 kg/ms and the free flow rate is less than about 70 minutes per 500 ml through standard enteral tubing. Monounsaturated fatty acids provide about 29% of energy, polyunsaturated fatty acids provide about 6% of energy, and saturated fatty acids provide about 5% of energy.

[0058] It will be understood that numerous modifications may be made to the specific embodiments described above without departing from the scope of the invention as defined in the claims.

Claims

1. A nutritional composition for diabetic patients, the composition comprising a protein source providing about 10% to about 20% of energy, a lipid source providing about 30% to about 50% of energy, a carbohydrate source providing about 35% to about 55% of energy, and a fibre mixture including a viscous soluble fibre and inulin, a hydrolysate of inulin, or both.

EP 0 898 900 A2

2. A liquid nutritional composition for diabetic patients, the composition comprising a protein source, a lipid source, a carbohydrate source, and a fibre mixture including a viscous soluble fibre and inulin, a hydrolysate of inulin, or both.
3. A nutritional composition according to claim 2 which has a viscosity, when measured at room temperature, of 0.015 to 0.03 kg/ms.
4. A powdered nutritional composition for diabetic patients, the composition comprising a protein source, a lipid source, a carbohydrate source, and a fibre mixture including a viscous soluble fibre and inulin, a hydrolysate of inulin, or both; the powdered nutritional composition being reconstitutable in an aqueous liquid for providing a liquid nutritional composition.
5. A nutritional composition according to any of claims 1 to 4 in which the fibre mixture further includes a source of insoluble dietary fibre.
6. A nutritional composition according to claim 5 in which the source of insoluble dietary fibre is pea hull fibre.
7. A nutritional composition according to claim 5 or claim 6 in which the ratio of soluble fibre, including inulin, to insoluble fibre is 1:3 to 3:1.
8. A nutritional composition according to any of claims 1 to 4 in which the fibre mixture comprises about 0.5 g/100 ml inulin, about 0.5 g/100 ml pectin or gum arabic, and about 0.5 g/100 ml pea hull fibre.
9. A nutritional composition according to any of claims 1 to 8 in which the lipid source includes monounsaturated fatty acids which provide about 25% to about 35% of the energy of the composition.
10. A nutritional composition according to claim 9 in which the lipid source further includes saturated fatty acids providing less than 10% of the energy of the composition.





(19)

Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 898 900 A3

(12)

EUROPEAN PATENT APPLICATION

(88) Date of publication A3:
26.04.2000 Bulletin 2000/17

(51) Int. Cl.⁷: A23L 1/308, A23L 1/0528,
A23L 1/305, A23L 1/30,
A23L 1/09

(43) Date of publication A2:
03.03.1999 Bulletin 1999/09

(21) Application number: 98201018.3

(22) Date of filing: 31.03.1998

(84) Designated Contracting States:
AT BE CH DE DK ES FI FR GB GR IE IT LI LU MC
NL PT SE
Designated Extension States:
AL LT LV MK RO SI

(72) Inventors:
• Jaussan, Véronique
1110 Morges (CH)
• Roessle, Claudia
1110 Morges (CH)
• Schweizer, Thomas
1052 Le Mont-Sur-Lausanne (CH)
• Bourguignon, Michel
14740 Lasson (FR)

(30) Priority: 23.06.1997 EP 97201915

(71) Applicant:
SOCIETE DES PRODUITS NESTLE S.A.
1800 Vevey (CH)

(54) Composition and method for providing nutrition to diabetics

(57) A nutritional composition for diabetics. The composition contains a protein source, a lipid source, and a carbohydrate source. The composition includes a fibre mixture containing a viscous soluble fibre and inulin, a hydrolysate of inulin, or both. The fibre mixture may also contain insoluble fibre. When in liquid form, the nutritional composition has a low viscosity and excellent tube flow rates.

EP 0 898 900 A3

European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 98 20 1018

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
X	EP 0 768 043 A (BRISTOL-MYERS SQUIBB) 16 April 1997 (1997-04-16) * page 3, line 36-41 * * page 4, line 39-43 * * claims 1-3, 11-13; example 4 *	1, 2, 5	A23L1/308 A23L1/0528 A23L1/305 A23L1/30 A23L1/09
A	WO 94 12541 A (RAFFINERIE TIRLEMONTTOISE) 9 June 1994 (1994-06-09) * page 15, line 3-33 * * page 1, line 32 - page 2, line 2 * * page 4, line 2-13 * * page 5, line 6-12 * * page 7, line 19-25; claims 1, 3, 12, 13, 15, 16, 19, 20, 27 *	1, 5	
A	FR 2 673 360 A (A.R.D.) 4 September 1992 (1992-09-04) * page 3, line 18-25 * * page 10, line 28 - page 11, line 2; claims 1-8; table 4 *	1, 5	
A	EP 0 692 252 A (RAFFINERIE TIRLEMONTTOISE) 17 January 1996 (1996-01-17) * page 7, line 29-35; claims *	1	TECHNICAL FIELDS SEARCHED (Int.Cl.6) A23L A61K
A	WO 91 13076 A (RAFFINERIE TIRLEMONTTOISE) 5 September 1991 (1991-09-05) * page 2, line 24-29 * * page 3, line 8-20 * * page 8, line 16-20; claims 1, 7, 12-14, 16, 19, 24, 25 *	1	
D, A	EP 0 504 055 A (ROUSSEL-UCLAF) 16 September 1992 (1992-09-16) * the whole document * & US 5 292 723 A	1	
-/-			
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 2 March 2000	Examiner Van Moer, A
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

EPO FORM 1800 (04/01)



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 98 20 1018

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (INCLUD)
A	EP 0 756 828 A (NUTRICIA) 5 February 1997 (1997-02-05) * the whole document *	1	
A	EP 0 627 490 A (SÜDZUCKER) 7 December 1994 (1994-12-07) * page 2, column 17-19 * * page 2, line 27-31; claims 1,10,11 *	1	
A	EP 0 745 333 A (CLINTEC NUTRITION) 4 December 1996 (1996-12-04) * page 2, line 41 - page 3, line 1 * * page 3, line 11-13 * * page 3, line 31-33 * * page 3, line 52-55 * * claims 1-4,7,8 *	1,2,4,5	
A	EP 0 265 772 A (ABBOTT LABORATORIES) 4 May 1988 (1988-05-04) * page 3, line 23-25 * * page 3, line 38-40 * * page 3, line 46-49 * * claim 16 *	1,2,5,9	
The present search report has been drawn up for all claims			TECHNICAL FIELDS SEARCHED (INCLUD)
Place of search THE HAGUE		Date of completion of the search 2 March 2000	Examiner Van Moer, A
CATEGORY OF CITED DOCUMENTS		T: theory or principle underlying the invention E: earlier patent document, but published on, or after the filing date D: document cited in the application L: document cited for other reasons A: technological background O: non-written disclosure P: intermediate document	
X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure P: intermediate document		A: member of the same patent family, corresponding document	

EPO FORM 1500 (04/01) (PAG 01)

ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.

EP 98 20 1018

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

02-03-2000

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 768043	A	16-04-1997	AU 703090 B	18-03-1999
			AU 6818896 A	24-04-1997
			CA 2187394 A	17-04-1997
			JP 9168374 A	30-06-1997
			SG 45509 A	16-01-1998
			US 5776887 A	07-07-1998
WO 9412541	A	09-06-1994	BE 1006377 A	09-08-1994
			AU 5459694 A	22-06-1994
			DE 69320982 D	15-10-1998
			DE 69320982 T	08-04-1999
			EP 0670850 A	13-09-1995
			ES 2122218 T	16-12-1998
			US 5660872 A	26-08-1997
FR 2673360	A	04-09-1992	NONE	
EP 692252	A	17-01-1996	US 5721345 A	24-02-1998
WO 9113076	A	05-09-1991	BE 1003826 A	23-06-1992
			AT 105295 T	15-05-1994
			AU 7235491 A	18-09-1991
			CA 2076647 A	24-08-1991
			DE 69101914 D	09-06-1994
			DE 69101914 T	18-08-1994
			EP 0515456 A	02-12-1992
			ES 2053322 T	16-07-1994
			JP 5504568 T	15-07-1993
			US 5659028 A	19-08-1997
EP 504055	A	16-09-1992	FR 2673812 A	18-09-1992
			AT 128606 T	15-10-1995
			CA 2062920 A	14-09-1992
			DE 69205192 D	09-11-1995
			DE 69205192 T	23-05-1996
			ES 2081066 T	16-02-1996
			JP 6169723 A	21-06-1994
			US 5292723 A	08-03-1994
EP 756828	A	05-02-1997	AU 702989 B	11-03-1999
			AU 6087196 A	06-02-1997
			DE 69506095 D	24-12-1998
			DE 69506095 T	24-06-1999
			ES 2123903 T	16-01-1999
			US 5792754 A	11-08-1998

EPO JOURNAL

For more details about this annex: see Official Journal of the European Patent Office, No. 12/82

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 98 20 1018

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

02-03-2000

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 627490 A	07-12-1994	DE 4316425 A	24-11-1994
		AT 150794 T	15-04-1997
		DE 59402195 D	30-04-1997
		DK 627490 T	08-12-1997
		ES 2102093 T	16-07-1997
		JP 7051083 A	28-02-1995
		US 5478732 A	26-12-1995
EP 745333 A	04-12-1996	AU 5072196 A	12-12-1996
		CA 2173780 A	02-12-1996
		JP 8322509 A	10-12-1996
		US 5904948 A	18-05-1999
EP 265772 A	04-05-1988	AT 83616 T	15-01-1993
		AU 602016 B	27-09-1990
		AU 8012987 A	28-04-1988
		CA 1330900 A	26-07-1994
		DE 3783206 A	04-02-1993
		ES 2053497 T	01-08-1994
		GR 3006667 T	30-06-1993
		HK 1003158 A	16-10-1998
		JP 2599400 B	09-04-1997
		JP 63123354 A	27-05-1988
		KR 9603888 B	23-03-1996
		US 4921877 A	01-05-1990

EPC form 100a

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

